

# MODELS FOR X-11 AND X-11 FORECAST PROCEDURES FOR PRELIMINARY AND REVISED SEASONAL ADJUSTMENTS

Kenneth F. Wallis  
University of Warwick

## 1. INTRODUCTION

Procedures for the seasonal adjustment of economic time series have typically been evaluated by studying their effect on a sample of actual time series. Recent proposals for amendments and extensions to existing methods have also been evaluated in the same way. Perhaps this approach is thought to be inevitable given that "there seems to be no ideal process of evaluating a method of adjustment" (Granger 1978, p. 55). In contrast, however, this paper continues a line of research in which the properties of the procedures themselves are studied, in the abstract. It is hoped that this will improve our general understanding of the performance of the existing methods and their extensions, and help to explain the results of the previous empirical studies.

The particular procedure considered is the Bureau of the Census Method II, Variant X-11 (Shiskin et al. 1967), which is widely used and is generally held to give satisfactory results in the seasonal adjustment of historical data. Our analysis proceeds by linear filter methods. The basic framework of a set of "time-varying" linear filters is presented by Wallis (1982), and further properties of these filters and their components are considered in this paper. The use of linear methods implies that attention is restricted to the performance of X-11 in additive mode (in which seasonal components are estimated as average differences from, not average ratios to, the trend-cycle), neglecting the option of graduating extreme irregular values.

The problem of first adjusting the current month's observation and then revising the seasonally adjusted figure as time goes by and more data become available has received much attention recently. A specific suggestion is that the subsequent revisions in the adjusted values might be reduced by applying X-11 not to the observed series alone, but to a series augmented by forecasts of future values. This has been put forward in association with various forecasting methods (compare Dagum 1975, 1979; Geweke 1978; Kenny and Durbin 1982); but since in all these methods the forecasts are calculated as linear combinations of observed values, our analysis can also encompass these X-11 FORECAST procedures. Many statistical agencies run the X-11 program only once a year, and at that time, project seasonal factors for the adjustment of the next 12 months' data to be used as the data become available. Kenny and Durbin (1982) and Wallis (1982) have argued that this practice

should be replaced by running the program every month, and it is not considered in this paper.

The linear filter representation of the X-11 seasonal adjustment procedure is presented in section 2. The filters range from the purely one-sided moving average implicit in the preliminary adjustment of the most recent observation, through a number of asymmetric moving averages, to the symmetric moving average implicit in the adjustment of historical data. The relations with forecast-augmented procedures are considered in section 3, and following a result given by Geweke (1978) and Pierce (1980), we define the property of internal consistency of a set of linear filters. The polynomial regression origins of smoothing filters are considered in section 4, and it is shown that whereas sets of filters constructed in this way are internally consistent, the Henderson moving averages used in X-11, for which a new interpretation is given, are not. In section 5, forecast-augmented procedures are used to provide models for which the X-11 procedure already minimizes the mean squared error of revisions, thus illuminating the empirical comparison of various methods. Particular attention is paid to important work by Kenny and Durbin (1982) and Dagum (1975, 1979). The relation to optimal signal extraction methods is briefly considered in section 6.

## 2. THE X-11 LINEAR FILTERS

The original observable monthly series, denoted by  $x_t$ , is the input to a linear filtering operation, of which the output is denoted by  $y_t$ . In the present context  $y$  is the seasonally adjusted value of  $x$ , but the basic framework can be applied to any problem of signal extraction, interpolation, extrapolation, and smoothing by linear filter methods. The X-11 program comprises a sequence of moving average or linear filter operations, but their net effect can be represented by a single set of moving averages. For a date sufficiently far in the past, the *final* or *historical* adjusted value  $y_t^{(m)}$  is obtained by application of the symmetric filter  $a_m(L)$

*The computational assistance and helpful comments of Peter Burridge are gratefully acknowledged. This research was supported by a grant from the Social Science Research Council.*

$$y_t^{(m)} = a_m(L)x_t = \sum_{j=-m}^m a_{m,j}x_{t-j}$$

where  $L$  is the lag operator and  $a_{m,j} = a_{m,-j}$ . This filter is described variously as a  $2m + 1$ -term moving average or as a symmetric filter of half-length  $m$ . For current and recent data, this filter cannot be applied, and truncated asymmetric filters are employed:

$$y_t^{(i)} = a_i(L)x_t = \sum_{j=-i}^m a_{i,j}x_{t-j}$$

For the filter  $a_i(L)$ ,  $i = 0, 1, \dots, m$ , the subscript  $i$  indicates the number of "future" values of  $x$  entering the moving average, that is, the number of negative powers of  $L$  that appear, or the negative of the lower limit of summation in the expression  $\sum a_{i,j}x_{t-j}$ . Equivalently, the superscript on  $y$  indicates that  $y_t^{(i)}$  is the adjusted value of  $x_t$  calculated from observations at times  $t - m, t - m + 1, \dots, t, \dots, t + i$ , and  $y_t^{(0)}$  is the *first-announced* or *preliminary* seasonally adjusted figure. For the X-11 filters considered here, the value of  $m$  is 84. (It is assumed that at the stage at which the program chooses a 9-, 13-, or 23-term Henderson moving average to estimate the trend-cycle, the 13-term average is chosen.) Thus the X-11 program can be represented as a set of 85 linear filters with, respectively, 0, 1,  $\dots$ , 84 coefficients of future values. There is no attempt to compensate for the truncation of the weights applied to future data by increasing the number of past values entering the moving average; thus, with few exceptions, all filters involve 84 past values, as indicated by the constant upper limit of summation in the above equation, although the remote weights are very small. In figures 1, 2, and 3, the weights and transfer functions of three filters of particular interest,  $a_0(L)$ ,  $a_{12}(L)$ , and  $a_{84}(L)$ , are reproduced from Wallis (1982), where details of their calculation can be found; the symmetric filter is also given in Wallis (1974).

At times  $t + i$  and  $t + k$ , two seasonally adjusted values  $y_t^{(i)}$  and  $y_t^{(k)}$  can be calculated, corresponding to the unadjusted value  $x_t$ , and the *revision* is defined as

$$r_t^{(i,k)} = y_t^{(k)} - y_t^{(i)}, 0 \leq i < k \leq m$$

This reflects the information in the "new" data  $x_{t+i+1}, x_{t+i+2}, \dots, x_{t+k}$ . The *total revision* from a given point in time  $t + i$  is  $r_t^{(i,m)}$ . Often statistical agencies run seasonal adjustment programs only once a year, so that the revisions then made are  $r_t^{(i,i+12)}$ . The first annual revisions  $r_t^{(0,12)}$  are of particular interest below.

### 3. FORECAST-AUGMENTED PROCEDURES AND THE INTERNAL CONSISTENCY OF SETS OF LINEAR FILTERS

In the X-11 FORECAST procedures, current data are adjusted not by the one-sided filter  $a_0(L)$  but by later filters applied to a series extended by forecasts. In practice, 12

such forecasts are used; thus, the procedures calculate the preliminary adjusted value not as  $a_0(L)x_t$  but as  $a_{12}(L)\tilde{x}_t$ , where the tilde ( $\tilde{\phantom{x}}$ ) indicates that the input to the filter is the augmented series  $x_1, \dots, x_t, \tilde{x}_{t+1}, \dots, \tilde{x}_{t+12}$ . Since the forecasts are calculated as linear combinations of observed values, the result is still a one-sided filter of the original data. Writing the forecasts as

$$\tilde{x}_{t+k} = \sum_{j=0}^{\ell} f_{kj}x_{t-j}, k = 1, \dots, 12$$







the new one-sided moving average is



$$\begin{aligned} \tilde{a}_0(L)x_t &= \sum_{k=1}^{12} a_{12,-k}\tilde{x}_{t+k} + \sum_{j=0}^m a_{12,j}x_{t-j} \\ &= \sum_{j=0}^{\max(m,\ell)} (a_{12,j} + \sum_{k=1}^{12} a_{12,-k}f_{kj})x_{t-j} \end{aligned}$$

Dagum (1975, 1979) bases the forecasts on ARIMA models fitted to the observed series, while Kenny and Durbin (1982) use autoregressive models fitted by stepwise regression methods to the first differences of the series.

Support for X-11 FORECAST procedures is provided by a result given by Geweke (1978) and Pierce (1980), namely that the asymmetric filter for which the total revision has smallest mean square is given by the application of the symmetric filter  $a_m(L)$  to a series extended to the extent necessary by *optimal* linear forecasts. Similarly the one-sided filter for which the first annual revision  $r_t^{(0,12)}$  has smallest mean square is  $\tilde{a}_0(L)$  above, provided that the forecasts  $\tilde{x}_{t+k}$  are optimal. (Many of the empirical evaluations compare the *total* revisions of various procedures, even though only 12 forecasts are employed in the X-11 FORECAST procedures; we return to this below.) In Geweke's words, "the best linear forecast of any given linear combination of  $x_t$ 's is the given linear combination of the forecasted  $x_t$ 's."

Given a symmetric moving average, these results provide a way of constructing asymmetric moving averages for the adjustment of recent data, based on optimal forecasting equations for a given  $x$ -process. We say that a set of linear filters constructed in this way is internally consistent, with respect to the given  $x$ -process. Likewise, a given set of linear filters can be examined for its internal consistency by asking what forecast function and, hence, what  $x$ -process is implied if  $a_i(L)$  is equivalent to the application of  $a_m(L)$  to a series extended by  $m - i$  forecasts, and is this  $x$ -process the same for all possible pairs. By considering two intermediate adjusted values  $y_t^{(i)}$  and  $y_t^{(k)}$ ,  $0 \leq i < k < m$ , and repeatedly applying Geweke and Pierce's result, in effect repeatedly taking conditional expectations, we see that an internally consistent set of filters minimizes revisions throughout the whole sequence of adjustments. Thus, if the filters  $a_i(L)$  and  $a_k(L)$  minimize the mean square of the revisions  $r_t^{(i,m)}$  and  $r_t^{(k,m)}$ , respectively, being identical to the application of  $a_m(L)$  to a sequence of observations augmented by  $m - i$  and  $m - k$  forecasts, respectively, then  $a_i(L)$  also minimizes the mean square of the revisions  $r_t^{(i,k)}$ ,





and is identical to the application of  $a_k(L)$  to a series extended by  $k - i$  forecasts. Note that while the theoretical result refers to an optimal forecast given the infinite past, the practical filters involve a finite past; hence, some restriction on the  $x$ -processes considered is necessary in order for the equivalences among filters to hold exactly. Finite-order autoregressive models clearly allow correspondences among the practical filters, but moving average models whose infinite autoregressive representations do not converge quickly allow only an approximation to an optimal forecast equivalence.

As an illustration, we examine the 5-term Henderson moving averages used for estimation of the trend-cycle component in the quarterly version of the X-11 program (Shiskin et al. 1967, app. B, table 3A):

$$\begin{aligned} a_0(L)x_t &= 0.670x_t + 0.403x_{t-1} - 0.073x_{t-2} \\ a_1(L)x_t &= 0.257x_{t+1} + 0.522x_t + 0.294x_{t-1} - 0.073x_{t-2} \\ a_2(L)x_t &= -0.073x_{t+2} + 0.294x_{t+1} + 0.558x_t \\ &\quad + 0.294x_{t-1} - 0.073x_{t-2} \end{aligned}$$

We seek the  $x$ -process and associated forecast coefficients implied by the assumption that the one-sided filter  $a_0(L)x_t$  is identical to  $a_2(L)\hat{x}_t$ , that is, the symmetric filter applied to the forecast-augmented series  $x_{t-2}, x_{t-1}, x_t, \hat{x}_{t+1}, \hat{x}_{t+2}$ . For an exact equivalence, the forecasts can only involve three observed  $x$ -values, so we postulate a third-order autoregressive process, with forecasts given by

$$\begin{aligned} \hat{x}_{t+1} &= \phi_1x_t + \phi_2x_{t-1} + \phi_3x_{t-2}, \\ \hat{x}_{t+2} &= \phi_1\hat{x}_{t+1} + \phi_2x_t + \phi_3x_{t-1} \\ &= (\phi_1^2 + \phi_2)x_t + (\phi_1\phi_2 + \phi_3)x_{t-1} + \phi_1\phi_3x_{t-2} \end{aligned}$$

Equating the coefficients of, respectively,  $x_t, x_{t-1}$  and  $x_{t-2}$  in  $a_0(L)x_t$  and  $a_2(L)\hat{x}_t$  then gives three polynomial equations for the  $\phi$ 's, which have four solutions. Writing them in the form  $\phi(L) = 1 - \phi_1L - \phi_2L^2 - \phi_3L^3$ , these are

$$\begin{aligned} (1 - 0.568L - 0.432L^2) \\ (1 - 3.027L - 1.493L^2) \\ (1 - 4.459L + 3.459L^2) \\ (1 - 4.027L + 2.534L^2 + 1.493L^3) \end{aligned}$$

Note that this mathematical problem has multiple solutions even when one is attempting to recover an autoregressive operator that has actually been used to construct the one-sided filter from the symmetric filter. One hopes, in general, that (at least) one of these solutions has a plausible statistical interpretation. To consider this it is more informative to express  $\phi(L)$  in terms of its roots  $\phi(L) = \Pi (1 - \lambda_i L)$ , whereupon the solutions can be written

$$\begin{aligned} (1 - L)(1 + 0.432L) \\ (1 + 0.432L)(1 - 3.459L) \\ (1 - L)(1 - 3.459L) \\ (1 - L)(1 + 0.432L)(1 - 3.459L) \end{aligned}$$

Of these, the first is acceptable as an ARIMA (1,1,0) model, but the remainder are not acceptable from a statistical point of view due to the presence of the explosive root. Thus, we conclude that  $a_0(L)$  is equivalent to the application of  $a_2(L)$  to a sequence of observations augmented by optimal forecasts for series obeying the first model, that is, for such a series  $a_0(L)$  minimizes the mean square of the revisions  $r_t^{(0,2)}$ .

The problem of nonlinearity does not arise when comparing  $a_0(L)$  with  $a_1(L)$ , or  $a_1(L)$  with  $a_2(L)$ , since only one forecast is involved, whose coefficients are (linear in) the  $\phi$ 's. Simple calculation gives the model relating  $a_0(L)$  to  $a_1(L)$  as

$$(1 - L)(1 + 0.424L)x_t = \epsilon_t$$

and the model relating  $a_1(L)$  to  $a_2(L)$  as

$$(1 - L)(1 + 0.493L)x_t = \epsilon_t$$

Although the three models are very similar, their coefficient values do not exactly coincide, and we conclude that these filters are not internally consistent in this sense.

#### 4. INTERNAL CONSISTENCY OF REGRESSION-BASED FILTERS

##### 4.1 Least squares polynomial regression

Filters used to estimate the trend-cycle component prior to estimation of the seasonal component are usually designed to reproduce, locally, a polynomial of prescribed degree  $d$ . A least squares estimate of the trend value then emerges as a linear function of the data, whose coefficients are independent of time, and these are the moving average coefficients. (See, for example, Kendall 1973, ch. 3.) Given a sample of  $n$  observations  $x_1, \dots, x_n$  placed in a column vector  $\mathbf{x}$ , and an  $n \times (d + 1)$  matrix  $\mathbf{Z}$  whose  $j$ th column comprises the  $(j - 1)$ th powers of the integers  $1, \dots, n$ , the trend estimate at time  $t$  is

$$y_t = \mathbf{z}'_t (\mathbf{Z}'\mathbf{Z})^{-1} \mathbf{Z}'\mathbf{x}$$

where  $\mathbf{z}'_t = (1, t, t^2, \dots, t^d)$ . Writing the coefficients of the observations in a column vector  $\mathbf{a}$ , with the coefficients of the earliest data entering first, then

$$y_t = \mathbf{a}'\mathbf{x}, \quad \mathbf{a} = \mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{z}_t$$

The coefficients are independent of time in the sense that, on indexing the data  $t_0 + 1, t_0 + 2, \dots, t_0 + n$  and using these integers to construct  $\mathbf{Z}$  and  $\mathbf{z}_t$  conformably, the same coefficients are obtained whatever the value of  $t_0$ . So when  $n = 2m + 1$ , it is conventional to let the time index run from  $-m$  to  $m$  for arithmetical convenience. (Similarly, the use of orthogonal polynomials simplifies the arithmetic without affecting the results.) If  $n$  is odd and  $t$  is the midpoint of the sample, then a symmetric moving average is obtained. Kendall (1973, app. A) presents Cowden's tabulations of moving averages for trend estimation at  $t = 0, 1, \dots, n, n + 1$ , for given values of  $n$  and  $d$ . Such a table does not correspond to a set of filters in the sense of section 2, but such a set is easily constructed by putting  $t = m + 1$  and considering in turn  $n = m + 1, m + 2, \dots, 2m + 1$ , and this set is internally consistent with respect to the regression model.

To see this, consider the filter applicable to  $n \leq 2m + 1$  data points and that applicable to  $k < n$  (but  $k > d$ ) data points. For internal consistency, the latter must be equivalent to the application of the former to the series  $x_1, \dots, x_k, \hat{x}_{k+1}, \dots, \hat{x}_n$ , where the forecasts are now obtained by polynomial regression. We partition the vector  $\mathbf{x}$  and matrix  $\mathbf{Z}$  into their first  $k$  and last  $(n - k)$  rows, thus

$$\mathbf{x}' = (\mathbf{x}'_1; \hat{\mathbf{x}}'_2), \quad \mathbf{Z}' = (\mathbf{Z}'_1; \mathbf{Z}'_2)$$

The forecasts  $\hat{x}_2$  obtained from polynomial regression based

on the data  $\mathbf{x}_1$  are

$$\hat{x}_2 = \mathbf{Z}_2(\mathbf{Z}'_1\mathbf{Z}_1)^{-1}\mathbf{Z}'_1\mathbf{x}_1$$

and applying the  $n$ -term moving average to the series  $\bar{\mathbf{x}}' = (\mathbf{x}'_1; \hat{\mathbf{x}}'_2)$  gives the trend estimate at time  $t$  as

$$\begin{aligned} y_t &= \mathbf{z}'_t (\mathbf{Z}'\mathbf{Z})^{-1} \mathbf{Z}'\bar{\mathbf{x}} \\ &= \mathbf{z}'_t (\mathbf{Z}'_1\mathbf{Z}_1 + \mathbf{Z}'_2\mathbf{Z}_2)^{-1} (\mathbf{Z}'_1\mathbf{x}_1 + \mathbf{Z}'_2\mathbf{Z}_2(\mathbf{Z}'_1\mathbf{Z}_1)^{-1}\mathbf{Z}'_1\mathbf{x}_1) \\ &= \mathbf{z}'_t (\mathbf{Z}'_1\mathbf{Z}_1)^{-1} \mathbf{Z}'_1\mathbf{x}_1 \end{aligned}$$

which is identical to the application of the corresponding  $k$ -term filter to the data  $\mathbf{x}_1$ .

##### 4.2 The Henderson moving averages

These filters are designed to reproduce a cubic polynomial trend. In obtaining a general expression for the weights of the symmetric filter, Henderson (1916) showed that three alternative smoothing criteria lead to the same result:

- (i) minimization of the variance of the third differences of the output series,
- (ii) minimization of the sum of squares of the third differences of the moving average coefficients,
- (iii) fitting a cubic by weighted least squares, with the sum of squares of third differences of the weights a minimum.

A more accessible presentation of the general derivation and a demonstration of the equivalence of these criteria is in the appendix of Kenny and Durbin (1982). A further interpretation can be obtained using the approach of Hannan (1970, pp. 186-7), as follows.

Augmenting the moving average coefficient vector  $\mathbf{a}$  with zeros to accommodate the differencing operations, criterion (ii), the sum of squares of the third differences of these coefficients, can be written

$$S = \mathbf{a}'\mathbf{V}\mathbf{a}$$

where  $\mathbf{V}$  is a symmetric band matrix with elements on the main diagonal and first three sub-diagonals equal to 20, -15, 6, and -1, respectively, all other elements being zero. Hannan shows that minimization of this general expression subject to the condition that the filter reproduces an arbitrary polynomial of degree  $d$  yields the solution

$$y_t = \mathbf{a}'\mathbf{x} = \mathbf{z}'_t (\mathbf{Z}'\mathbf{V}^{-1}\mathbf{Z})^{-1} \mathbf{Z}'\mathbf{V}^{-1}\mathbf{x}$$

In Hannan's formulation,  $\mathbf{V}$  is the covariance matrix of the input series, and he notes that, as is well known, this result corresponds to using the best linear unbiased regression of the input series on the polynomials to estimate the trend value at the relevant time point. In this case, the particular matrix  $\mathbf{V}$  is the covariance matrix of the moving average process

$$u_t = (1 - L)^3 \varepsilon_t$$

and so we have a further interpretation of the Henderson filters, namely as generalized least squares polynomial regression subject to this error structure. If  $n$  is odd and  $t$  is the midpoint of the sample, then the coefficients are again symmetric, and using this approach, we can readily verify their published values.

By setting  $t = m + 1$  and considering in turn  $n = m + 1, m + 2, \dots$ , and  $2m + 1$ , the above expression gives a set of filters of the kind discussed earlier. This set is again internally consistent with respect to the present regression model. The proof of this is a tedious generalization of the proof for the least squares case given in section 4.1, taking account of the need to define  $\hat{x}_{k+1}, \dots, \hat{x}_n$  as the best linear unbiased predictors of  $x_{k+1}, \dots, x_n$  given  $x_1, \dots, x_k$ .

On computing sets of filters for  $m = 4, 6$ , and  $11$  in this way, we immediately notice that, except for the symmetric filter, the coefficients differ from those used in X-11. The source of the asymmetric moving averages given by Shiskin et al. (1967, app. B, tables 3B-3D) corresponding to the symmetric 9-, 13-, and 23-term Henderson moving averages is something of a mystery: They are not due to Henderson (1916), no alternative source is cited, and they are not the result of applying Henderson's criteria to the design of asymmetric filters. Our calculations in general find that the weights given to current data are somewhat larger than those in the tabulated filters used in X-11. Kenny and Durbin (1982) report that the "one-sided version is constructed essentially by assuming that the series to be smoothed can be extended by a straight line fitted by least squares," which is their own empirical finding for the monthly filters. For the quarterly 5-term filters analysed in section 3, we do not claim that the autoregressive models we have deduced were actually used to construct the asymmetric filters, but we note that the models imply that the forecast is approximately given as the average of the last two observed values, which may be a more plausible interpretation of the actual derivation. Basing asymmetric filters on forecasts obtained from a different model than that embodied in the symmetric filter leads to internal inconsistency. In turn, this leads one to expect that improvements over the existing one-sided seasonal adjustment filters could be made for series well-described by the model underlying the symmetric filter and, hence, satisfactorily adjusted by it.

## 5. FORECASTING INTERPRETATIONS OF THE RELATIONS BETWEEN SELECTED FILTERS

### 5.1 Autoregressive models relating $a_0(L)$ to $a_{12}(L)$

The filters under consideration require a finite amount of past data, and in section 3 the possibility was suggested of finding an autoregressive model and associated forecast functions such that an existing filter  $a_i(L)$  is already equivalent to the application of  $a_k(L)$  to a series augmented by  $k - i$  forecasts. For such a model, the mean square of the revision  $r_t^{(i,k)}$  is already at a minimum. Much research

has focussed on the adjustment of current data, and the practical X-11 FORECAST procedures augment the observed series with 12 forecasts, and so first we consider the relation between the X-11 filters  $a_0(L)$  and  $a_{12}(L)$ .

To obtain a general characterization, we follow the approach of Kenny and Durbin (1982), in which forecasts are obtained from autoregressive models. They estimate these by regressing the first difference  $\Delta x_t$  on  $\Delta x_{t-1}, \dots, \Delta x_{t-25}$ , reducing the set of regressors by the Efronson stepwise method. Neglecting the restriction implied by the use of the first difference operator, the general model is

$$\phi(L)x_t = \epsilon_t$$

where  $\phi(L)$  is of degree 26. For this model we construct the forecast coefficients for 1-step, 2-step,  $\dots$ , 12-step forecasts

$$\hat{x}_{t+k} = \sum_{j=0}^{25} f_{kj}x_{t-j}, \quad k=1, \dots, 12$$

by starting with the expression for the one-step forecast

$$\hat{x}_{t+1} = \phi_1 x_t + \phi_2 x_{t-1} + \dots + \phi_{26} x_{t-25}$$

and then using, sequentially, the expression for the  $k$ -step forecast

$$\begin{aligned} \hat{x}_{t+k} &= \phi_1 \hat{x}_{t+k-1} + \dots + \phi_{k-1} \hat{x}_{t+1} \\ &+ \phi_k x_t + \dots + \phi_{26} x_{t+k-26} \quad k=2, \dots, 12 \end{aligned}$$

We then seek the  $\phi$ 's that equate the one-sided filter implied by the application of  $a_{12}(L)$  to the forecast-augmented series, namely

$$\bar{a}_0(L) = \sum_{j=0}^{25} \left[ a_{12,j} + \sum_{k=1}^{12} f_{kj} a_{12,-k} \right] L^j + \sum_{j=26}^m a_{12,j} L^j$$

to  $a_0(L)$ . Note that the use of a finite autoregression implies that the later coefficients in the new filter are simply those of  $a_{12}(L)$ , and so the new filter cannot be exactly equated to  $a_0(L)$ . (Comparing the coefficients of  $a_0(L)$  and  $a_{12}(L)$  plotted in figure 1, we see that there is little difference in the coefficients at lags greater than 25 except at lag 36.) Equating the earlier coefficients in the two filters gives

$$a_{0,j} = a_{12,j} + \sum_{k=1}^{12} f_{kj} a_{12,-k} \quad j = 0, 1, \dots, 25$$

a set of 26 nonlinear equations in the 26 unknown  $\phi$ 's which we solve by numerical methods. (A nonlinear least squares algorithm is used: at a solution the residual sum of squares is zero.) As in the simple example of section 3, there are multiple solutions to this numerical problem, and while we cannot claim to have examined them all, indeed we are not sure how many there are, the model given below has the most plausible statistical interpretation and is also suggested by the models relating successive pairs of filters.

$i$	$\phi_i$	$i$	$\phi_i$
1	-.60	14	.27
2	-.40	15	.14
3	-.26	16	-.15
4	.13	17	.15
5	-.19	18	-.20
6	.38	19	.23
7	-.16	20	-.16
8	.29	21	.15
9	-.13	22	-.08
10	.12	23	-.09
11	.11	24	-.39
12	-.28	25	.59
13	-.21	26	.01

To interpret this general autoregression, we calculate the roots of the autoregressive operator,  $\phi(L) = \prod(1 - \lambda_i L)$ . There are 2 real roots,  $(1 + 0.982L)$  and  $(1 + 0.018L)$ , and 12 complex conjugate pairs, whose modulus and argument (in radians) are as follows:

mod	arg	mod	arg
1.033	0.10	0.981	0.27
0.958	0.51	0.945	0.82
0.983	1.06	0.965	1.37
0.985	1.57	0.975	1.89
0.982	2.10	0.978	2.42
0.985	2.62	0.988	2.81

Noting that the arguments in the first block of six pairs of roots are approximately  $k\pi/6$ ,  $k = 0, 1, \dots, 5$ , these together with the first real root can be approximated as  $(1 - L)(1 - L^{12})$ , the differencing operator commonly employed in ARIMA models for seasonal time series. While this solution contains a complex conjugate pair of roots with modulus greater than one, it is one of the least explosive solutions to this problem we have found, and no solution representing a stationary autoregression has been observed. In empirical time series analysis, the use of as many as 26 coefficients to specify a model would be thought excessive, and likewise the relations between the filters may be more readily interpreted in terms of seasonal ARIMA models parameterized more parsimoniously, as considered in section 5.2.

The implication of this result is that, for a series with such autoregressive structure, the existing one-sided X-11 filter  $a_0(L)$  minimizes the mean square of the first annual revision  $r_t^{(0,12)}$ , and no X-11 FORECAST procedure of this kind will lead to an improvement. Of course for series having substantially different autoregressive structures, the appropriate X-11 FORECAST procedure will outperform X-11. Attention is then drawn to Kenny and Durbin's anomalous finding that for one group of nine series (those with MCD = 1) the X-11  $a_0(L)$  gives better results than the X-11 FORECAST  $\bar{a}_0(L)$ . If this finding is not due to sampling variation, then three possible explanations are suggested.

The first possibility, on which we have little further to say, is that the various nonlinearities in the practical procedures vitiate our linear analysis. The second is that the comparison is based as in other studies (e.g., Kuiper 1978), on the total revision  $r_t^{(0,m)}$  of the two methods and not the first annual revision  $r_t^{(0,12)}$ . The one-sided filter that minimizes the mean square of  $r_t^{(0,m)}$  is  $a_m(L)$  applied to a series augmented by  $m$  forecasts; call this  $a_0^*(L)$ . On the other hand, the filter being considered is  $\bar{a}_0(L)$ , namely  $a_{12}(L)$  applied to a series augmented by 12 forecasts. If the filters were internally consistent, then  $a_{12}(L)$  would be equivalent to the application of  $a_m(L)$  to a series augmented by  $m - 12$  forecasts and  $a_0(L)$  could be obtained either by applying this  $a_{12}(L)$  to a series augmented by 12 forecasts or as  $a_0^*(L)$ . However, our earlier results suggest and later results confirm that the X-11 filters are not internally consistent, and we have the possibility that for these series the X-11  $a_0(L)$  is a better approximation to  $a_0^*(L)$  than is  $\bar{a}_0(L)$ , hence the result.

A third explanation involves the possible nonoptimality of the autoregressive forecast, which might apply if the appropriate models for the series had important moving average components. However, the form of the seasonal ARIMA models identified by Kenny and Durbin for these series is little different in this respect from those for the series for which X-11 FORECAST methods dominate, and without knowing the coefficient values, this possibility cannot be assessed. Using the conventional Box-Jenkins notation  $(p, d, q) \times (P, D, Q)_{12}$ , the nine series giving the anomalous results have  $p + d + D = 3$ , and this distinguishes them from the other series. Five of the nine models are  $(0, 2, 1) \times (0, 1, 1)_{12}$ , and the remaining four also have  $P = 0$  and  $Q = 1$ . Leaving aside questions about the existence of measures such as MCD for models of this kind (Burridge 1982) and despite the fact that if MCD=1, the X-11 program applies the 9-term Henderson moving average to estimate the trend-cycle component, not the 13-term average used in our linear filter calculations, the results suggest that such models might emerge when we consider equivalences between filters based on ARIMA models.

## 5.2 ARIMA models relating $a_0(L)$ , $a_{12}(L)$ and $a_{34}(L)$

In this section, we report the results of a comparable exercise to that of the previous section, except that seasonal ARIMA models replace the finite autoregressive model. That is, for a given seasonal ARIMA model, we construct the forecast coefficients and, hence, the X-11 FORECAST filter coefficients

$$\bar{a}_{0,j} = a_{12,j} + \sum_{k=1}^{12} f_{kj} a_{12,-k}, \quad j = 0, 1, \dots, m,$$

and then seek the model and its associated parameter values such that  $\bar{a}_0(L)$  is as close as possible to  $a_0(L)$ . For a given model, the parameter values are chosen to minimize the unweighted sum of squared deviations between the filter

coefficients. The choice among candidate models is partly based on the sum of squares, but we also have in mind the various criteria used to interpret such models when they are fitted to observed time series. For comparative purposes, we include in the search the three models available in the automatic version of Statistics Canada X-11 ARIMA (Dagum 1979), and it turns out that one of these is our preferred model. The results for these three models are shown below. The final column can be compared to the "zero-forecast" sum of squares

$$\sum_{j=0}^m (a_{0,j} - a_{12,j})^2 = 0.0477$$

We note that the autoregressive operator in the preferred model has a root close to unity, but setting it equal to unity worsens the fit, and we retain the  $(2, 1, 2) \times (0, 1, 1)_{12}$  model for further comparisons below, where this feature is less pronounced.

Again the interpretation is that for a series obeying the model  $(2, 1, 2) \times (0, 1, 1)_{12}$ , with the above parameter values, the existing X-11 filter  $a_0(L)$  is a close approximation to the X-11 ARIMA filter that minimizes the mean square of the first annual revision  $r_t^{(0,12)}$ . In empirical comparisons of the two methods, the advantage of X-11 ARIMA will increase as the  $x$ -process differs more extensively from that given above. In practice, we note that substantial variation can be observed as parameter values change within a given model.

To consider filters that minimize the mean square of the total revision  $r_t^{(0,m)}$ , we repeat this exercise for the filter

$a_0^*(L)$  obtained by applying the symmetric filter  $a_m(L)$  to a series augmented by  $m$  forecasts:

$$a_{0,j}^* = a_{mj} + \sum_{k=1}^m f_{kj} a_{mk}, \quad j = 0, 1, \dots, m$$

The preferred model is again  $(2, 1, 2) \times (0, 1, 1)_{12}$ , and the results for the three Dagum models are shown below.

$$\sum_{j=0}^m (a_{0,j} - a_{mj})^2 = 0.0696$$

Results in earlier sections cast doubt on the internal consistency of the X-11 filters, and while the models relating these three filters are not identical, the form of the preferred model is the same in each case, and the coefficients are remarkably similar. To complete the picture, we report the coefficient values for the  $(2, 1, 2) \times (0, 1, 1)_{12}$  model that best approximates  $a_{12}(L)$  by the application of  $a_m(L)$  to a series augmented by  $m - 12$  forecasts:

$$\begin{aligned} (1 - 1.08L + 0.21L^2)(1 - L)(1 - L^{12})x_t \\ = (1 - 1.42L + 0.70L^2) \\ (1 - 0.60L^{12})\epsilon_t \end{aligned}$$

Once more these values are similar, suggesting that whatever adjusted values empirical comparisons are based on,  $y_t^{(0)}$ ,  $y_t^{(12)}$ , or  $y_t^{(m)}$ , then for series well-described by this model, X-11 will produce approximately the same results as X-11 FORECAST methods.

<i>Estimated model relating <math>a_0(L)</math> to <math>a_{12}(L)</math></i>	<i>Sum of squared deviations</i>
$(1 - 1.37L + 0.39L^2)(1 - L)(1 - L^{12})x_t = (1 - 1.49L + 0.76L^2)(1 - 0.69L^{12})\epsilon_t$	0.0096
$(1 - L)^2(1 - L^{12})x_t = (1 - 1.32L + 0.63L^2)(1 - 0.74L^{12})\epsilon_t$	0.0111
$(1 - L)(1 - L^{12})x_t = (1 - 0.25L)(1 - 0.69L^{12})\epsilon_t$	0.0188
<i>Estimated model relating <math>a_0(L)</math> to <math>a_m(L)</math></i>	<i>Sum of squared deviations</i>
$(1 - 1.23L + 0.35L^2)(1 - L)(1 - L^{12})x_t = (1 - 1.45L + 0.73L^2)(1 - 0.62L^{12})\epsilon_t$	0.0114
$(1 - L)^2(1 - L^{12})x_t = (1 - 1.33L + 0.58L^2)(1 - 0.65L^{12})\epsilon_t$	0.0124
$(1 - L)(1 - L^{12})x_t = (1 - 0.21L)(1 - 0.66L^{12})\epsilon_t$	0.0218

## 6. OPTIMAL SIGNAL EXTRACTION

The problem of seasonal adjustment is formulated as a problem of optimal signal extraction by Grether and Nerlove (1970) and solved using methods presented by Whittle (1963). It is assumed that the observed series  $x_t$  is the sum of two unobserved uncorrelated stationary components,  $s_t$  and  $u_t$ , respectively the seasonal and nonseasonal com-

ponents. Assuming that the stochastic structure of  $s_t$  and  $u_t$  and hence that of  $x_t$ , is known Whittle gives expressions for  $s_t^{(k)}$ , the minimum mean squared error estimate of  $s_t$  given data  $\{x_\tau; \tau \leq t + k\}$ . These expressions are linear filters of the data, in general, of semi-infinite extent; the case  $k = 0$  gives a symmetric filter. Extensions of these results to models in which the components follow nonstationary ARIMA schemes are given by Cleveland and Tiao (197



(see also Pierce, 1979), who present a components model for which the X-11 symmetric filter  $a_m(L)$  approximates the optimal signal extraction filter. The relations between the filters considered in the previous section are examined in the light of such models by Burridge and Wallis (1982).

The practical difficulty in implementing seasonal adjustment methods based on optimal signal extraction results is that of identifying models for the components from the observed series, given that different components models can lead to the same overall model. Burman (1980) makes some practical proposals in this respect, and implements a seasonal adjustment method based on the resulting decomposition. We are not yet able to incorporate this approach into our comparisons, but anticipate that for series well-described by the models for which the X-11 filters approximate optimal signal extraction filters, little benefit from the new methods will be found in empirical comparisons.

#### REFERENCES

- BURMAN, J. P. (1980), "Seasonal Adjustment by Signal Extraction," *Journal of the Royal Statistical Society, A*, 143, 321-337.
- BURRIDGE, P. (1982), Contribution to the discussion of Kenny and Durbin (1982).
- BURRIDGE, P., and WALLIS, K. F. (1982), "Unobserved-Components Models for Seasonal Adjustment Filters," unpublished paper, Department of Economics, University of Warwick.
- CLEVELAND, W. P., and TIAO, G. C. (1976), "Decomposition of Seasonal Time Series: A Model for the Census X-11 Program," *Journal of the American Statistical Association*, 71, 581-587.
- DAGUM, E. B. (1975), "Seasonal Factor Forecasts From ARIMA Models," *Bulletin of the International Statistical Institute*, 46, pt. 3, 203-216.
- \_\_\_\_\_ (1979), "The X-11 ARIMA Seasonal Adjustment Method: Outline of the Methodology," catalogue 12-564E, Statistics Canada.
- GEWEKE, J. (1978), "The Revision of Seasonally Adjusted Time Series," *Proceedings of the Business and Economic Statistics Section, American Statistical Association*, pp. 320-325.
- GRANGER, C. W. J. (1978), "Seasonality: Causation, Interpretation, and Implications" (with discussion), in *Seasonal Analysis of Economic Time Series*, A. Zellner, ed., pp. 33-55, Washington, D.C.: Bureau of the Census.
- GRETHER, D. M., and NERLOVE, M. (1970), "Some Properties of 'Optimal' Seasonal Adjustment," *Econometrica*, 38, 682-703.
- HANNAN, E. J. (1970), *Multiple Time Series*, New York: Wiley.
- HENDERSON, R. (1916), "Note on Graduation by Adjusted Average," *Transactions of the Actuarial Society of America*, 17, 43-48.
- KENDALL, M. G. (1973), *Time Series*, London: Griffin.
- KENNY, P. B., and DURBIN, J. (1982), "Local Trend Estimation and Seasonal Adjustment of Economic and Social Time Series" (with discussion), *Journal of the Royal Statistical Society, A*, 145, 1-41.
- KUIPER, J. (1978), "A Survey and Comparative Analysis of Various Methods of Seasonal Adjustment," in *Seasonal Analysis of Economic Time Series*, A. Zellner, ed., pp. 59-76, Washington, D.C.: Bureau of the Census.
- PIERCE, D. A. (1979), "Signal Extraction Error in Nonstationary Time Series," *Annals of Statistics*, 7, 1303-1320.
- \_\_\_\_\_ (1980), "Data Revisions With Moving Average Seasonal Adjustment Procedures," *Journal of Econometrics*, 14, 95-114.
- SHISKIN, J.; YOUNG, A. H.; and MUSGRAVE, J. C. (1967), *The X-11 Variant of the Census Method II Seasonal Adjustment Program*, Technical Paper 15 (rev.), Washington, D.C.: Bureau of the Census.
- WALLIS, K. F. (1974), "Seasonal Adjustment and Relations Between Variables," *Journal of the American Statistical Association*, 69, 18-31.
- \_\_\_\_\_ (1982), "Seasonal Adjustment and Revision of Current Data: Linear Filters for the X-11 Method," *Journal of the Royal Statistical Society, A*, 145, 76-85.
- WHITTLE, P. (1963), *Prediction and Regulation by Linear Least-Square Methods*, London: English Universities Press.

# COMMENTS ON "MODELS FOR X-11 AND X-11 FORECAST PROCEDURES FOR PRELIMINARY AND REVISED SEASONAL ADJUSTMENTS" BY KENNETH F. WALLIS

John Geweke  
University of Wisconsin-Madison

The revision problem in the seasonal adjustment of time series has long been familiar to those responsible for the production of seasonally adjusted series, and somewhat more recently has drawn the attention of academics as well. Analytical treatment of the problem has centered on the fact that if seasonal adjustment procedures constitute the application of a linear filter, then the mean square error of revision will be minimized if the filter is applied to the available data augmented by optimal linear forecasts of the unavailable data. Exploitation of this fact to reduce actual revisions in published seasonally adjusted time series is made difficult by the need to parameterize time series and then estimate the parameters before forecasts can be produced, to say nothing of the political and bureaucratic problems involved in the routine application of procedures which require subjective judgments by highly trained practitioners.

In this paper, Kenneth Wallis has provided models which can be used to study further the degree to which official procedures conform to the "forecasting paradigm." Attention is focused on Census X-11, which (at least in very similar variants) is the basis for most seasonal adjustment throughout the world. Census X-11 does not explicitly forecast unavailable data, but uses special filters near the beginnings and ends of series when the number of data points required is not available. Wallis compares these filters and asks whether they can be interpreted as the application of the historical Census X-11 filter to the series augmented by linear forecasts. The filters are said to be internally consistent if there exists a linear forecast such that the application of historical Census X-11 to the augmented series results in the set of filters employed. If the forecast turns out to be the best linear forecast of a given series, then the set of filters is optimally internally consistent with respect to the series. I found this method of analysis very useful. It provides a quick answer to the internal consistency question, and allows us to organize some of the empirical results that have appeared in recent years. In what follows, I shall first elaborate on the paper's finding that Census X-11 is internally inconsistent, and then indicate how the method of analysis developed here might be further employed to make sense of empirical results.

Before turning to the analysis, I shall present some evidence in support of a stylized fact which has been obscured by the method of presentation in many empirical papers (including Kénnny and Durbin 1981, the most recent and accessible of the papers cited by Wallis): The magnitude of

the revision of seasonally adjusted data is much more a function of the series in question than it is of the method used to cope with the revision problem. In table 1, Census X-11 is compared with "optimal" procedures, with forecasts made in four alternative ways: Using a canonical factorization of the estimated spectral density (X-11S), a 3-year autoregression (X-11A), an ARIMA model (0, 1, 1) × (0, 1, 1) (X-11 ARIMA (1)), and an ARIMA model (0, 2, 2) × (0, 1, 1) (X-11 ARIMA (2)). The variation in revisions across series is much greater than the variation in revisions across methods (including Census X-11), and differences among the four alternative procedures are small compared with the difference between them and Census X-11.

The analytical work in Wallis's paper shows that Census X-11 is internally inconsistent. In view of the fact that the filters used near the beginnings and ends of series were not constructed in any optimal way—at least to the best of anyone's recollection—this is not surprising. It is interesting to evaluate the implications of the discrepancy for mean square revision. To keep matters simple, consider the five-term Henderson moving average cited as illustration in the paper. It is shown that the forecasting model implicit in the filters for this average is of the form

$$\Delta x_t = b \Delta x_{t-1} + \epsilon_t$$

However, the value of  $b$  implicit depends on which filters are being compared; there are three filters involved, and the pair-wise comparisons yield the values of  $-0.432$ ,  $-0.424$ , and  $-0.493$  for  $b$ . The five-term Henderson moving average may be written

$$-0.073\Delta x_{t+2} + .221\Delta x_{t+1} + .779x_t + .294x_{t-1} - .073x_{t-2}$$

The revision between time  $t$  and time  $t + 2$  is

$$-0.073\Delta x_{t+2} + .221\Delta x_{t+1} + (.073b^2 - .221b)\Delta x_t$$

*John Geweke is Associate Professor in the Department of Economics, 1180 Observatory Drive, University of Wisconsin, Madison, Wis., 53706. National Science Foundation grant SES8005606 provided partial financial support.*

If the process were

$$\Delta x_t = c \Delta x_{t-1} + \epsilon_t$$

then the mean square revision would be

$$\text{MSR}(b,c) = (.073b^2 - .221b)(.073b^2 - .221b + .442c - .146c^2) - .032266c + .054170$$

Suppose that in fact  $c = -.432$ . If  $b = -.432$ ,  $\text{MSR} = .05620708$ . If  $b = -.424$ , then  $\text{MSR} = .05621222$ , an increase of .0046 percent in root mean square revision; and if  $b = -.493$ , it is 0.5651684, an increase of .2752 percent. This seems to suggest that the internal inconsistency of Census X-11—at least in the five-term Henderson moving average—is not a practical problem.

This finding is driven by the fact that, locally, there is no return to optimal forecasting because

$$\partial \text{MSR}(b,c) / \partial b |_{b=c} = 0 \tag{1}$$

Given a change in  $c$ , it makes no difference (locally!) whether the old forecasting method is used or the newer optimal one is employed:

$$\begin{aligned} \partial \text{MSR}(b,c) / \partial c |_{b=c} &= d \text{MSR}(c,c) / dc \tag{2} \\ &= -.021316c^3 + .096798c^2 - .097682c - .0322166 \end{aligned}$$

Since this expression is generally nonzero, variation in  $c$  may well lead to a reduction in MSR even when the forecasting procedure is not modified. This kind of result is suggested by (1), but (2) seems strong; whether the latter result holds more generally warrants investigation.

This analysis suggests that there ought not to be much variation in the revision of seasonally adjusted series for different sensible forecasting procedures, but that substantial variation will arise across different series for the same procedure. For nonsensical forecasting procedures, this need not be the case, and the analogue of this fact in the context of the analytical examples employed here is shown in table 1: Globally,  $b$  and  $c$  are about equally important in explaining variation in MSR. This analysis further appears to model well the results reported in table 2.

TABLE 1. EVALUATION OF MSR ( $b,c$ )

c	b				
	-0.8	-0.4	0.	0.4	0.8
-0.8	0.030	0.045	0.080	0.120	0.155
-0.4	.072	.057	.067	.088	.110
0.	.104	.064	.054	.060	.071
0.4	.126	.067	.041	.035	.038
0.8	.136	.064	.028	.014	.011

These considerations lead to a fourth explanation (beyond the three suggested by Wallis) for the Kenny-Durbin (1981) finding that for a particular group of series, the conventional Census X-11 program produces smaller revisions on average than do various forecasting procedures. The series in question were those with the lowest months of cyclical dominance (MCD), i.e., those with the shortest span over which average absolute changes in the trend component exceed absolute changes in the irregular component. For smoother series, a fixed process which extrapolates trend values will predict better than it will for more erratic series. For such series we may interpret Census X-11 as providing a sensible set of forecasts. Since the X-11 forecasting weights have no variance (unlike the weights in procedures requiring estimation), it seems plausible that mean square revision in X-11 could well be smaller than that in procedures with more flexible forecasting capabilities. This interpretation is supported by Kenny and Durbin's finding that the latter procedures compare increasingly favorably with conventional Census X-11 as the MCD of the series increase.

REFERENCES

GEWEKE, JOHN (1978), "The Revision of Seasonally Adjusted Time Series," *Proceedings of the Business and Economic Section, American Statistical Association*, 320-325.

\_\_\_\_\_ (1980), "An Evaluation and Comparison of X-11, X-11S, X-11A, and X-11 ARIMA," University of Wisconsin memorandum.

KENNY, P. B., and DURBIN, J. (1981), "Local Trend Estimation and Seasonal Adjustment of Economic and Social Time Series," *Journal of the Royal Statistical Society A*, forthcoming.

TABLE 2. ROOT MEAN SQUARE PROPORTIONAL REVISION OF RELATIVE CHANGES

Series	X-11	X-11S	X-11S	X-11 ARIMA(1)	X-11 ARIMA(2)
1	1.96	1.35 (31.1%)	1.35 (31.1%)	1.32 (32.6%)	1.33 (32.1%)
2	.96	.70 (27.1%)	.73 (24.0%)	.74 (22.9%)	.75 (21.9%)
3	.85	.67 (21.1%)	.67 (21.1%)	.67 (21.1%)	.68 (20.0%)
4	14.01	10.17 (27.4%)	10.30 (26.5%)	10.43 (25.6%)	11.01 (21.4%)
5	10.34	7.62 (26.3%)	7.61 (26.4%)	6.99 (32.4%)	6.98 (32.5%)
6	4.55	3.11 (31.6%)	2.99 (34.3%)	3.88 (14.7%)	3.80 (16.5%)

Source: Geweke (1980). These series are (1) general merchandise retail sales; (2) motor vehicles and automotive equipment inventories; (3) inventories of groceries and related products; (4) one-family housing starts, northeast; (5) motor vehicle assembly operations; (6) sales by blast furnaces and steel mills. For details on computations, see Geweke (1978). Figures in parenthesis indicate improvement over X-11.

**COMMENTS ON "MODELS FOR X-11 AND X-11 FORECAST PROCEDURES  
FOR PRELIMINARY AND REVISED SEASONAL ADJUSTMENTS" BY  
KENNETH F. WALLIS**

Adi Raveh

Kenneth Wallis studies procedures for seasonal adjustment of economic time series in the abstract, rather than trying to evaluate their effects on sample time series, as is usually done. This interesting and constructive paper continues his time-varying linear filter method (Wallis 1974, 1981). In an ingenious and concise presentation the author provides tools to check whether linear filters are internally consistent. Among other findings, it is shown that sets of filters constructed by polynomial regression are internally consistent and that X-11 filters (for additive model) are not. Practically, X-11 filters do not deviate terribly from being internally consistent.

Comparisons between X-11 and X-11 FORECAST for various approximations to models are given in the paper as well. For series well-described by X-11 ARIMA based on the  $(2, 1, 2) (0, 1, 1)_{12}$  model with specific parameter values, empirical comparisons suggested that X-11 will produce approximately the same results as X-11 FORECAST methods. Hence, Wallis' paper contributes a lot to our general understanding of the performance of X-11 and X-11 FORECAST.

The problem of revision of the seasonal adjustment of empirical time series may be decomposed into mainly two parts: (a) revision due to structure of the data, namely the specific series in question and (b) revision due to the method used to deal with the problem. This paper focused on the latter part as is required by its title. On the other hand, it seems that most of the magnitude of the revision is essentially a function of the first part as already mentioned by Geweke (1980).

Consider an ideal series which has fixed seasonality and no irregularity. Then the periodicity-free series known as seasonally adjusted data need not be revised at all. The variation in revisions across methods for such series, as well as series that have an irregular component and moving seasonality superimposed gradually, can also be a plausible basis for comparison purposes.

The use of the linear filter methods implies that attention is restricted to the additive model and neglects the option of graduating extreme irregular values. Most of the empirical time series which are analyzed by X-11 (90 percent or more) or other decomposition methods are analyzed using multiplicative models. Thus, from the application point of view, perhaps efforts should be devoted to improving our general understanding of the multiplicative model and the mixed multiplicative-additive model in further research (Raveh

1981). Young (1968, p. 456) indicated that. "The effect of the nonlinear operations is not very large and for some purposes can be ignored." Nevertheless, it is worthwhile to assess more precisely the revisions obtained by nonlinear operations. Wallis' paper has obviously made great progress in studying some theoretical properties of X-11. However, further research has to pay attention to the option of graduating extreme values which are often used as well as the nonlinear filters.

Another point which the paper did not take into consideration is the constraints for seasonal factors that are built into X-11. These constraints force the seasonal factors of a calendar year to add up to zero when applying the additive model and add up to 12 (or 1200 percent) for the multiplicative version. On account of these constraints, different shifts of the series backward or forward yield different estimation results for the main components: Trend, seasonal, and irregular. For example, let a given series cover  $m$  whole years from January through December. Then, shift the series 6 months ahead, or in other words, January is called July, February is called August, etc. Now the series includes  $(m - 1)$  whole years plus 2 half years. Apply X-11 to the shifted series as well as to the original one. It yields different results. For example, the seasonal factors for January of the original series are not the same as those for July of the shifted series. The above facts indicate that different revisions are obtained for the same series shifted backward or forward. In other words, the revision is affected by the location of the last observation within the year.

Another question worth mentioning is whether it is necessary to revise seasonal adjustments at all for series that have a fixed or nearly fixed seasonal pattern. For such series it is plausible, sometimes, to use the seasonal pattern which has already been estimated to adjust current data. In order to be able to discriminate between fixed and moving seasonality, they have to be defined properly or as simply as possible.

To conclude my comments, I would like to present a family

*The author is on leave from Hebrew University, Jerusalem, and currently is an ASA Junior Research Fellow at the Bureau of the Census. This work was initiated when the author was associated with Stanford University.*

of coefficients for measuring the amount of linear association different from the classical product-moment coefficient. This family is based on ideas that quantify the magnitude of the slopes of a given series over time. In some sense, X-11 uses the measure proposed below for forecasting the seasonal factors 1 year ahead.

A series  $X_1, \dots, X_n$  is perfectly linear if and only if

$$X_i - X_{i-1} = X_{i-1} - X_{i-2} \quad \text{for all } i = 3, \dots, N \quad (1)$$

or

$$\Delta^2 X_i = \Delta X_i - \Delta X_{i-1} = 0 \quad i = 3, \dots, N \quad (2)$$

where  $\Delta X_i = X_i - X_{i-1}$ . On the other hand, for any three adjacent observations  $\Delta^2 X_i$  can be either positive or negative. The maximum of  $\Delta^2 X_i$  for any triple  $X_i, X_{i-1}, X_{i-2}$  is obtained when  $\Delta X_i = -\Delta X_{i-1}$  and its value is

$$\Delta^2 X_i = (\Delta X_i - (\Delta X_{i-1})) = 2\Delta X_i.$$

It easy to verify that the quantity

$$\frac{\sum_{i=3}^N |\Delta^2 X_i|^p}{\sum_{i=2}^N |\Delta X_i|^p} \quad (3)$$

varies between 0 and 1. Let us focus on one coefficient from a family based on the above quantity, say when  $p = 2$ . This measure is given as

$$0 \leq c = \text{LIN}(X) = \text{LIN}(X_1, \dots, X_n) = 1 - \frac{\sum_{i=3}^N (\Delta^2 X_i)^2}{4 \sum_{i=2}^N (\Delta X_i)^2} = 1 - K \leq 1 \quad (4)$$

$\text{LIN}(X) = c = 1$  ( $K = 0$ ) if and only if the series is perfectly linear. In other words, its slope remains constant over time.  $\text{LIN}(X) = c = 0$  ( $K = 1$ ) if and only if the slope of every two consecutive observations changes its sign but its absolute value remains the same. This means that in some sense a series like  $(a, b, a, b, \dots, b)$  where  $a \neq b$  is the most dissimilar to linear association. For both cases, when  $\text{LIN}(X) = 1$  or 0, perfect prediction is obtained.  $\text{LIN}(X) \approx 0$  if the series is locally linear, namely, there are very few turning points (relative to the length of the series) and in between the series is linear.

Given sets of filters are said to be internally consistent if there exists a linear forecast such that the application of the decomposition (e.g., X-11) method to the augmented series results in the set of filters employed. Let us suggest an

approach for forecasting, which is in some sense the "other way around" from what has been done in Wallis' paper. To be specific, we find forecasts for which the past and forecast series are consistent, namely, the same in the short-term or at least in the range of the required forecasting. The structures can be measured by appropriate coefficients of goodness-of-fit. The forecasted values are then the augmented series such that the goodness-of-fit (or structure) would be the same as for the original series.

Let us call the above principle the Persistent Structure Principle. The "structure" can be measured as linear or monotone or any other function of time, depending on the researcher's loss function. The very simple case of predicting one unit ahead based on linear trend is presented here. A more general case of monotone trend will be given in Raveh (1982).

Let us demonstrate our approach by using the coefficient  $c = \text{Lin}(X)$  given in (4) and based on the "ideal" linear conditions in (1). Our principle requires that the estimated forecast  $\hat{X}_{n+1}$  to be such that  $\text{LIN}(X, \hat{X}_{n+1})$  for the augmented series would be equal to the  $\text{LIN}(X)$  of the original series. Hence, by equating  $\text{LIN}(X, \hat{X}_{n+1}) = \text{LIN}(X)$ , the required  $\hat{X}_{n+1}$  is found to be

$$\hat{X}_{n+1} = aX_n + (1 - a)X_{n-1} \quad (5)$$

where  $a$  can be either  $a_1 = \frac{2\sqrt{K} - 2}{2\sqrt{K} - 1}$  or  $a_2 = \frac{2\sqrt{K} + 2}{2\sqrt{K} + 1}$  and  $K$  is defined in (4). Thus,  $\hat{X}_{n+1}$  is a combination of the two last observations. The weights  $a, 1 - a$  are functions of the entire series.

For the case of perfect linear correlation,  $\text{LIN}(X) = 1$  (or  $K = 0$ ) and  $a = 2$ , this yields the equation

$$\hat{X}_{n+1} = 2X_n - X_{n-1} = X_n + (X_n - X_{n-1})$$

In the other extreme case of perfect negative linear association,  $\text{LIN}(X) = 0$  (or  $K = 1$ ) and  $a = 0$ . Thus,

$$\hat{X}_{n+1} = X_{n-1}$$

The procedure in X-11 forecasts the seasonal factor for the  $j$ th month 1 year ahead, with fixed weights in the following way:

$$\hat{S}_{j,n+1} = S_{j,n} + 1/2[S_{j,n} - S_{j,n-1}] = 1.5S_{j,n} - 0.5S_{j,n-1} \quad (6)$$

In light of the above approach X-11 adopts  $a_2 = 1.5$  or  $K = 0.25$  or  $\text{LIN}(X) = 0.75$ . Thus, the forecasts of  $\hat{S}_{n+1}$  are in some sense based on rough approximations of linear relationships between  $S_1, S_2, \dots, S_n$ .

Equation (6) is the formula used for prediction purposes when the identified model is an ARIMA (1,1,0) as in (7):

$$(1 - \theta B)\Delta S_t = a_t \quad (7)$$

where  $B$  is back shift operator and  $a_t$  a white noise. The parameter  $\theta$  in (7) is equivalent to  $-(1-a)$  in (5). In other words, in order to predict seasonal factors 1 year ahead, X-11 uses an ARIMA (1,1,0) model with  $\theta = .5$  (or  $a = 1.5$ ) as a constant and not by means of an optimal parameter estimation process. Recently, we found that for many economic series from the Bureau of the Census data base the optimal (in terms of the above Persistent Structure Principle) coefficient  $a$  is very close to 1.5.

Needless to say, LIN(X) is based heavily on the last two observations. For forecasting purposes we recommend the use of more stable relations involving any four observations rather than any consecutive three observations. Thus, for example, stronger conditions for linearity are given in (8) below instead of the conditions in (1) and (2).

$$\frac{X_i - X_j}{i - j} = \frac{X_k - X_\ell}{k - \ell}$$

$$\text{and } n \geq i > j \geq 1 \text{ and } n \geq k > \ell \geq 1. \quad (8)$$

## REFERENCES

- GEWEKE, J. (1980), "An Evaluation and Comparison of X-11, X-11S, X-11A, and X-11 ARIMA," University of Wisconsin memorandum.
- RAVEH, A. (1981), "A Numerical Nonmetric Approach for Analyzing Time Series Data," *Communications in Statistics, Part A: Theory and Methods*, A10(8), 809-821.
- \_\_\_\_\_ (1982), Forecasting Economic Time Series Using the Persistent Structure Principle. Submitted for publication.
- WALLIS, K. F. (1974), Seasonal Adjustment and Relations Between Variables. *Journal of the American Statistical Association* 69, 18-31.
- \_\_\_\_\_ (1981), Seasonal Adjustment and Revisions of Current Data: Linear Filters for the X-11 Method. *Journal of the Royal Statistical Society A*, 144, forthcoming.
- YOUNG, A. H. (1968), Linear Approximations to the Census and BLS Seasonal Adjustment Methods. *Journal of the American Statistical Association* 63, 445-471.