



Mental Illness Among America's Youth: Results from the 2011 National Survey on Drug Use and Health

*Peter Delany, Director
Center for Behavioral Health Statistics and Quality
Substance Abuse and Mental Health Service
Administration*



March 2013






Overview: In 2011, 70,109 respondents participated in a computer assisted, self administered, face to face interview. Survey is representative nationally and in each State.

Results from the 2011 National Survey on Drug Use and Health:

- Youth aged 12 to 17 years continue to report substantial rates of major depressive episodes
- Among adults, young adults 18 to 25 years, have the highest reported rates of mental illness, suicidal thoughts, and major depressive episodes
- Co-occurring substance use and dependence is not uncommon among individuals who suffer from mental illness
- Females are likely to report emotional problems

Major Depressive Episode in the Past Year among Youths Aged 12 to 17, by Gender: 2004-2011


Percent with MDE in the Past Year



⁺ Difference between this estimate and the 2011 estimate is statistically significant at the .05 level.


Past Year Substance Dependence or Abuse among Youths Aged 12 to 17, by Major Depressive Episode in the Past Year: 2011

Percent Dependent or Abusing Substance




Past Year Mental Health Service Use among Youths Aged 12 to 17, by Gender: 2011

Percent Using Mental Health Services
in the Past Year



Any Mental Illness in the Past Year among Adults Aged 18 or Older, by Age: 2011


Percent with Any Mental Illness (AMI) in the Past Year



Age Group


Co-Occurring Mental Illness and Substance Use Disorder in the Past Year among Adults Aged 18 to 25, by Gender: 2011

Percent with Co-Occurring Mental Illness and Substance Use Disorder in the Past Year



Any Mental Illness in the Past Year among Adults Aged 18 or Older, by Age and Gender: 2011

Percent with Any Mental Illness (AMI) in the Past Year




Note: Male vs. Female is significant at the 0.05 level for every comparison within age group.

Note: All pairwise comparisons of age groups by gender are significant at the 0.05 level within gender. This does not include the 18 or Older category.

Serious Mental Illness in the Past Year among Adults Aged 18 or Older, by Age and Gender: 2011

Percent with Serious Mental Illness (SMI) in the Past Year




Note: Male vs. Female is significant at the 0.05 level for every comparison within age group.

Note: All pairwise comparisons of age groups by gender are significant at the 0.05 level within gender. This does not include the 18 or Older category.

Suicidal Thoughts in the Past Year among Adults Aged 18 or Older, by Age and Gender: 2011



Percent with Serious Thoughts of Suicide in the Past Year




Note: Male vs. Female is significant at the 0.05 level for every comparison within age group except for the 50 or Older group.


Note: All pairwise comparisons of age groups by gender are significant at the 0.05 level except 26 to 49 vs. 50 or Older among Males. This does not include the 18 or Older category.

Major Depressive Episode in the Past Year among Adults Aged 18 or Older, by Age and Gender: 2011

Percent with Major Depressive Episode (MDE) in the Past Year




Note: Male vs. Female is significant at the 0.05 level for every comparison within age group.
 Note: 18 to 25 vs. 50 or Older is significant at the 0.05 level within gender.
 Note: 26 to 49 vs. 50 or Older is significant at the 0.05 level within gender.



Past Year Mental Health Service Use among Adults Aged 18 or Older, by Type of Care: 2002-2011

Percent Using Mental Health Services in the Past Year



- ← Any Type of Care
- ← Prescription Medication
- ← Outpatient
- ← Inpatient



⁺ Difference between this estimate and the 2011 estimate is statistically significant at the .05 level.